


Problem Set #2


Due Wed, Apr 22th

1. [25 marks] Consider the PRR-planar mobile manipulator shown in the figure below. Assume that m_1 , m_2 , and m_3 are point masses.



- Find the Jacobian matrix in frame $\{0\}$, relating velocity at the end-effector to the joint velocities. Use $[x \ z \ \alpha]^T$ as the operational coordinates where $\alpha = \theta_2 + \theta_3$.
Hint: Use a geometry to find the Jacobian. In this case DH parameters might confuse you.
- Find the singularities. Explain the physical meaning.
- Derive the joint space kinetic energy matrix A and the gravity vector G .
Use $C_1 I_1 = C_2 I_2 = C_3 I_3 = 0$.
- Derive the operational space kinetic energy matrix Λ and the gravity vector P at the end-effector.
- Consider the configuration when $\theta_2 = 45^\circ$ and $\theta_3 = 90^\circ$. Assume $d_1 = l_2 = l_3 = 1m$ and $m_1 = m_2 = m_3 = 1kg$ and the manipulator is at rest.
Hint: At rest, centrifugal and coriolis force is zero.
 - Find the joint torques required to compensate for gravity.
 - Determine the joint torques required to accelerate the base at $1m/s^2$ while maintaining the joint position of the arm. Use joint space equations of motion. Discuss your results.
 - Determine the joint torques required to accelerate the end-effector at $1m/s^2$ along the z_0 -axis. Use operational space equations of motion. Discuss your results.


2. [25 marks] *Operational space* Consider the PRR-spatial manipulator shown in the figure below. Assume that m_4 is a point mass, $m_1 = 0$, and $I_i = 0$ for all joints. The task of this manipulator is to *position* the end-effector in the 3-D workspace. The configuration shown below is when $q_2 = 0$.



- Select a set of operational space coordinates for this task.
- Find the Jacobian matrix corresponding to the operational coordinates, in frame $\{0\}$.
- Find the Jacobian matrix corresponding to the operational coordinates, in frame $\{3\}$.
- Determine the kinematic singularities. Draw each one and explain its physical meaning.
- Find the joint space kinetic energy matrix $A(\mathbf{q})$ and the joint space gravity vector $G(\mathbf{q})$.
- Find the configuration where the kinetic energy of the manipulator is minimum, for a given $\dot{\mathbf{q}}$ (you may assume $\dot{\mathbf{q}} \neq \mathbf{0}$).
- Take $m_2 = m_3 = 1kg$, and $L_3 = 1m$. Find the operational space kinetic energy matrix ${}^3\Lambda(\mathbf{q})$ and the operational space gravity vector ${}^3P(\mathbf{q})$ in frame $\{3\}$.

3. [25 marks] *Kinematics, Dynamics and Instantaneous Inverse Kinematics*

Consider the PRR planar manipulator shown in the figure below. The task of this manipulator is to *position* the end-effector. You are also given that $m_1 = m_2 = 1$, $L_1 = L_2 = 1$, $m_3 = 0$ and $I_{zz1} = I_{zz2} = 1$.



- Choose operational space coordinates appropriate for the task.
- Is this manipulator redundant with respect to the task? If so, what is the degree of the redundancy?
- Find the Jacobian matrix.
- Are there any singularities for this manipulator? If so, find the configurations from the Jacobian matrix and describe the resulting restriction in motion.
- For the configuration with $q_2 = 0^\circ$ and $q_3 = -45^\circ$, find the operational space kinetic energy matrix Λ .

$$\text{Use } A = \begin{bmatrix} m_1 + m_2 & m_2 L_1 c_2 & 0 \\ m_2 L_1 c_2 & m_2 L_1^2 + I_{zz1} + I_{zz2} & I_{zz2} \\ 0 & I_{zz2} & I_{zz2} \end{bmatrix}$$

- Consider another configuration: $q_2 = 90^\circ$ and $q_3 = 0^\circ$. For a given δx , we would like to find the joint angles δq . Determine δq using the pseudo inverse and the dynamically consistent inverse of J separately.
- When $\delta x = [0.1 \ 0]^T$, calculate your δq for both cases and discuss the difference.
- What are the null spaces for both cases? Explain the possible motions without changing δx .